

File: ISIS-PG-FMT502_inpXX_dat.xml

Description of 'inp04.dat'; Randlsq Program A-priori Input File Format, where XX denotes the run-number ID for the best-fit solution from RAND, which was retained for that particular body. These run-number IDs are fortunately unique between each of the measured Saturnian moons such that XX: 03=Mimas, 04=Dione, 05=Iapetus, 06=Tethys, 07=Rhea, 14=Enceladus. Note, this input format is identical to the Randlsq program output file format.

Created as part of a project to put planetary geodesy control networks on the web. These control networks are from ISIS Planetary Geodesy Software (formerly RAND/USGS Planetary Geodesy (RUPG) Software).

Version: 2015.10.05

=====

Filename: inpXX.dat (example: inp04.dat)

Description: Primary input of a priori information, such as approximate positions for the body pole position and rotation rate, control point positions (and optionally their uncertainties), and camera position and orientation.

File format:

Group 1 - Control point locations ("npoi" records, note 2):

Phi	1-24	D24.16	Latitude of control point (degrees).
Lamda	25-48	D24.16	Longitude of control point (degrees).
If iew=0, then east longitude. If iew=1, then west longitude.			
Radius	49-72	D24.16	Radius of control point (km).
Pointid	73-78	A6	Point identification (unitless).

Sample (from inp04.dat):

=> 0.7848108923922052D+01 0.3546883340205007D+03 0.5600000000000000D+03
13<=

Group 2 - Camera orientation and position (4 x "npic" records, note 3):

Record 4-1:

JulianDate	1-24	D24.16	Julian date when picture was taken (days).
Imageid	25-36	A12	Image identification. Usually flight data sequence (FSC) or similar image number (unitless).

- 64-78 A15 "JULIAN_DATE&FDS".

Record 4-2:

s(i,1) 1-24 D24.16 X component of spacecraft position
vector in J2000.0 coordinates (km).

s(i,2) 25-48 D24.16 Y component of spacecraft position
vector in J2000.0 coordinates (km).

s(i,3) 49-72 D24.16 Z component of spacecraft position
vector in J2000.0 coordinates (km).

- 73-78 A6 "SXSYSZ".

Record 4-3:

c(i,1) 1-24 D24.16 J2000.0 right ascension of optical axis
of picture (degrees).

c(i,2) 25-48 D24.16 J2000.0 declination of optical axis of
picture (degrees).

c(i,3) 49-72 D24.16 Twist angle of picture (degrees).

- 73-78 A6 "C1C2C3".

Sample (from inp04.dat):

```
=> 0.2444556029881424D+07       3493011
JULIAN_DATE&FDS
0.6848283167042200D+06   0.7565547282075000D+04   -0.1227938257916000D+06
SXSYSZ
0.1805908868573520D+03   0.1010498000432080D+02   0.3308782231715768D+03
C1C2C3<=
```

Notes:

1. Currently read from randlsq program unit 12.
2. "npoi" is the number of control points. See the "Solution Parameterization" file (format "ISIS-PG-FMT531.doc") for input of this.
3. "npic" is the number of images. See the "Solution Parameterization" file (format "ISIS-PG-FMT531.doc") for input of this.
4. "PLANET" angles (records such as no. 4-4 above) and the use of the runstring option "PLANET=YES" should be used for the Moon and eventually other objects where the closed formulae in randlsq (subroutine iau2000.F) are not sufficiently accurate to represent the rotation of the body in question.
5. Lines beginning with a "#" will eventually be treated as comments.
6. Earlier versions of randlsq for non-lunar solutions used 5 character control point names, but all solutions now use 7 character names.

7. The input of uncertainties for control points was added to the operational version of randlsq after 2005 November 17. These uncertainties are used to weight the given parameters by addition to the diagonal elements of the normal matrix. Latitude and longitude uncertainties here are converted to radians. Latitude weights are then one over the latitude uncertainty squared. Longitude weights are one over (the longitude uncertainty divided by the absolute value of the cosine of the latitude) squared. Radius weights are one over the radius uncertainty squared. Note that in experimental and briefly operational (2005 October) versions before this date, the uncertainties were (incorrectly) not squared. So the entered uncertainties were actually treated as entered variances. Also note that if uncertainty values are specified here, they are used to weight the parameters, overriding the use of "RAND style" weights as set using the gweight or sweight values in the Parameter Input File. (And as a current bug, these changed weights are not yet reflected in the output of the RAND style weights at the top of the residual output file, under the heading "WEIGHTS".)

=====

Reference: Model, program, and format generally follow that specified in:

Colvin, Tim R. (1992). "Photogrammetric Algorithms and Software for Spacecraft Optical Imaging Systems," _ A RAND NOTE _, N-3330-JPL.

Note that the original format indicates the use of 5 character control point names.

=====

Document History:

Begun 2006.08.10 by B. Archinal, based on RUPG-FMT5012.doc.

Modifications: Modified 2015.10.05 by G. Cushing for web release.

=====

(End of document.)